arXiv:cond-mat/0511253v1 [cond-mat.other] 10 Nov 2005

On the stability of the coherent state of a two-level atom Bose gas in the resonant laser field at zero temperature

L. A. Maksimov, A. V. Paraskevov*

Kurchatov Institute, Kurchatov Sq. 1, Moscow 123182, Russia

It is shown that a Bose gas of two-level atoms in the intense resonant laser field at zero temperature is a mixture of two condensates with a definite ratio of the densities. The criteria of stability are found for the stationary states of such system against the increment of the amplitudes of quasi Bogoliubov elementary excitations. Besides the usual acoustic mode the gap mode is shown to exist and the magnitude of the gap is proportional to the laser field amplitude. The involvement of the gas nonideality under definite conditions results in an instability and decay of the condensates.

PACS: 03.75.Kk, 03.75.Mn

1. Introduction

After the experimental realization of the Bose-Einstein condensation in a gas [1] and later in a binary mixture of Bose gases [2] in magnetic traps at ultralow temperatures the adequate theoretical description of such systems is undoubtedly actual. This is confirmed by a great number of theoretical papers on the classification [3] of all states of a Bose mixture (within the Thomas-Fermi approximation) and its dynamics (longwave collective excitations [4], metastable states [5], the spatial separation of the mixture components [6]). In most papers the mixture is described by two Gross-Pitaevskii equations and the concrete distinction between the mixture components is not considered. The representation of the mixture components as atoms in the ground and excited states is employed in the works on studying the scattering [7] and absorption [8] of the laser radiation by the Bose condensate. However, the question about an existence of the stationary coherent state of the interacting system, "mixture + laser field", has not been considered yet. The present work is devoted to the theoretical investigation of this problem. In particular, the equilibrium ratio of the condensate densities is determined as a function of the resonant laser field amplitude and proximity to the saturation, the spectrum of quasi Bogoliubov elementary excitations is found, and the criteria of stability in the system of two condensates in the laser field are determined with the involvement of the recoil momentum and nonideality of a gas.

---

*e-mail: paraskevov@kurm.polyn.kiae.su

Let us consider a weakly non-ideal Bose gas of two-level atoms in the resonant laser field of the high intensity at zero temperature. The monochromatic laser field, acting on the Bose condensate of atoms in the ground state (for simplicity, we neglect noncondensate particles), converts some fraction of the atoms into the dipole-excited state so that each excited atom moves with the same velocity governed by the recoil momentum. If the laser field intensity is so high that one can neglect the spontaneous decay of excited atoms with respect to the induced emission, then, during the small time as compared with the time between collisions of excited atoms with quasiparticles of the condensate of unexcited atoms, the macroscopically large number of dipole-excited atoms moving with the same velocity is formed and the second Bose condensate appears. Below we do not treat the process of the formation of this condensate and assume that the state with two condensate has been already formed, taking as a condition that the natural linewidth of dipole transition is $\gamma \approx 0$. The criterion of a smallness for γ will be given in Sec. 2. At last, from the viewpoint of maximal simplicity of the effect treatment, the atomic system is considered as spatially homogeneous.

2. The ground state of two-level gas in the laser field

The Hamiltonian for the system consisting of N_a unexcited atoms and N_b dipole-excited atoms in the laser field ($V = 1, \hbar = 1$) reads

$$\begin{gathered} H_{tot} = H_{gas} + H_{int} + H_{ph},\ H_{gas} = H_a + H_b + H_{ab}, \\ H_a = \sum_p (\tfrac{p^2}{2m}) \hat{a}_p^+ \hat{a}_p + \sum_{p_1+p_2=p_3+p_4} \tfrac{1}{2} U_{aa} \hat{a}_{p_1}^+ \hat{a}_{p_2}^+ \hat{a}_{p_3} \hat{a}_{p_4}, \\ H_b = \sum_p (\tfrac{p^2}{2m} + \omega_0) \hat{b}_p^+ \hat{b}_p + \sum_{p_1+p_2=p_3+p_4} \tfrac{1}{2} U_{bb} \hat{b}_{p_1}^+ \hat{b}_{p_2}^+ \hat{b}_{p_3} \hat{b}_{p_4}, \\ H_{ab} = \sum_{p_1+p_2=p_3+p_4} U_{ab} \hat{a}_{p_1}^+ \hat{b}_{p_2}^+ \hat{b}_{p_3} \hat{a}_{p_4}. \end{gathered} \tag{1}$$

Here $\hat{a}_p$ and $\hat{b}_p$ are the atom annihilation operators with momentum p in the ground and dipole-excited states, respectively, and $U_{ik} = 4\pi a_{ik}/m$ are the parameters of the pair interaction expressed in terms of the corresponding scattering lengths. To exclude the problem of collapse, we suppose $U_{ik} > 0$. The interaction of the gas with a single-mode laser field is described by the Hamiltonian

$$H_{int} = \sum_p (g_k \hat{c}_k^+ \hat{a}_p^+ \hat{b}_{p+k} + g_k^* \hat{b}_{p+k}^+ \hat{a}_p \hat{c}_k), \tag{2}$$

where $\hat{c}_k$ and $\hat{c}_k^+$ are the absorption and production operators of a photon with wave vector $\vec{k}$ and frequency ω_k. For simplicity, the laser radiation is assumed to be linearly polarized in the direction of unit polarization vector $\vec{e}_k$. The atom-field coupling constant has a usual form

$$|g_k|^2 = 2\pi\omega_k \left|(\vec{e}_k, \vec{d})\right|^2$$

and is related to the natural linewidth of the excited level for a single atom via dipole matrix element d, $\gamma = \frac{4}{3}k_0^3 d^2$ where $k_0 = \frac{\omega_0}{c}$. To simplify the notation, the factor $\exp(i\vec{k}\vec{r})$ in the matrix element of the interaction between a single atom and laser field is omitted. Lastly, the Hamiltonian of free photons is given by

$$H_{ph} = \omega_k \hat{c}_k^+ \hat{c}_k.$$

The dynamics of the system is described by the equations

$$\begin{gathered} i\frac{\partial \hat{a}_p}{\partial t} = \frac{p^2}{2m}\hat{a}_p + \sum_{p+p_2=p_3+p_4} U_{aa}\hat{a}_{p_2}^+\hat{a}_{p_3}\hat{a}_{p_4} + \sum_{p_1+p=p_3+p_4} U_{ab}\hat{b}_{p_1}^+\hat{b}_{p_3}\hat{a}_{p_4} + g_k\hat{c}_k^+\hat{b}_{p+k}, \\ i\frac{\partial \hat{b}_p}{\partial t} = (\frac{p^2}{2m} + \omega_0)\hat{b}_p + \sum_{p+p_2=p_3+p_4} U_{bb}\hat{b}_{p_2}^+\hat{b}_{p_3}\hat{b}_{p_4} + \sum_{p_1+p=p_3+p_4} U_{ab}\hat{a}_{p_1}^+\hat{b}_{p_3}\hat{a}_{p_4} + g_k^*\hat{a}_{p-k}\hat{c}_k, \\ i\frac{\partial \hat{c}_k}{\partial t} = \omega_k\hat{c}_k + \sum_p g_k\hat{a}_p^+\hat{b}_{p+k}. \end{gathered} \tag{3}$$

Let us consider the system which consists of three coherent subsystems, namely, two condensates produced by atoms in the ground and dipole-excited states and the laser field. Let us find the ground state of this system, neglecting noncondensate particles. In what follows, it is more convenient to deal with usual complex quantities rather than with operators. Therefore the use of the representation of coherent states is appropriate. Let us determine a coherent state as

$$|a\rangle = \exp(\sum_p a_p\hat{a}_p^+)\,|0\rangle\,,$$

which can be considered as a direct product of all eigenvectors $|a_p\rangle$ so that $\hat{a}_p|a_p\rangle = a_p|a_p\rangle\,, \langle a_p|\,\hat{a}_p^+ = \langle a_p|\,a_p^*$ and concerning the states with different momenta. Vector ($|0\rangle$ denotes a direct product of all empty atom and photon states. For dipole-excited atoms and photons, the coherent states are determined similarly:

$$|b\rangle = \exp(\sum_p b_p\hat{b}_p^+)\,|0\rangle\,, \quad |c\rangle = \exp(c_k\hat{c}_k^+)\,|0\rangle\,.$$

Taking the recoil into account, condensates of atoms in the ground and dipole-excited states should move relatively each other. Let us denote the momenta of atoms belonging to these condensates $\vec{p}_a$ and $\vec{p}_b = \vec{p}_a + \vec{k}$, respectively. If the gas is at rest as a whole, $\vec{p}_a = -\vec{k}N_b/(N_a+N_b)$. Postmultiplying each of equations (3) by vector $|a\rangle\,|b\rangle\,|c\rangle$, we obtain the set of three equations for the complex fields a_{p_a}, b_{p_b}, c_k:

$$\begin{gathered} i\frac{\partial a_{p_a}}{\partial t} = (\frac{p_a^2}{2m} + U_{aa}N_a + U_{ab}N_b)a_{p_a} + g_k c_k^* b_{p_b}, \\ i\frac{\partial b_{p_b}}{\partial t} = (\frac{p_b^2}{2m} + \omega_0 + U_{bb}N_b + U_{ab}N_a)b_{p_b} + g_k^* a_{p_a} c_k, \\ i\frac{\partial c_k}{\partial t} = \omega_k c_k + g_k a_{p_a}^* b_{p_b}. \end{gathered} \tag{4}$$

Hereafter we consider a gas in the condensate state. Therefore N_a and N_b are macroscopically large quantities, i.e., $N_a,\ N_b \gg 1$. We seek a stationary solution in the form

$$a_{p_a} = ae^{-i\varepsilon_a t},\ b_{p_b} = be^{-i\varepsilon_b t},\ c_k = ce^{-i\omega t}. \tag{5}$$

Note that a stationary solution (5) takes place only under condition of the exact resonance between the renormalized laser frequency and the difference in energies of the excited and ground states of an atom

$$\omega = \varepsilon_b - \varepsilon_a. \tag{6}$$

From (4) we get

$$\begin{gathered} \varepsilon_a a = \Omega_a a + g_k c^* b, \\ \varepsilon_b b = \Omega_b b + g_k^* a c, \\ (\varepsilon_b - \varepsilon_a)\, c = \omega_k c + g_k a^* b. \end{gathered} \tag{7}$$

Here the following notations are introduced

$$\Omega_a = \frac{\vec{p}_a^{\,2}}{2m} + U_{aa}N_a + U_{ab}N_b,\ \Omega_b = \frac{\vec{p}_b^{\,2}}{2m} + \omega_0 + U_{ab}N_a + U_{bb}N_b,\ N_a = |a|^2,\ N_b = |b|^2.$$

Quantities ε_a, ε_b and N_a (or N_b since we consider that the given total gas density is $N = N_a + N_b = const$) are unknown. The third equation in (7) determines the renormalization of the laser frequency due to dipole transitions. This renormalization can be neglected if the laser field intensity is sufficiently high (formally, this means that $H_{ph} \gg H_{int}$, implying matrix elements) so that

$$N_c = |c|^2 \gg \frac{|g_k|^2 N_a N_b}{\omega_k^2}. \tag{8}$$

The condition of the smallness of the natural linewidth results directly from this inequality

$$\gamma \ll \omega_k \frac{N_c k_0^3}{N_a N_b}. \tag{9}$$

Inequality (8) is worthwhile to compare with the initially implied condition of the sufficiently high field intensity in order to neglect spontaneous decays with respect to the induced transitions. In the terms of the notations introduced this condition is given by

$$N_c \gg \frac{2}{\pi} k^3. \tag{10}$$

Note that, if one takes resonance approximation ($|\omega_k - \omega_0| \ll \omega_k, \omega_0$) into account, inequality (10) yields the condition known in the laser physics for saturation of the population of excited state, when an atom with the equal probability is in the ground and excited states so that $N_a = N_b = N/2$. The inequality is expressed quantitatively as

$$|\omega_k - \omega_0| \sim \gamma \ll |A| \,, \qquad (11)$$

where $|A| = |g_k| \sqrt{N_c}$ is the Rabi frequency in the conventional terminology. From (11) we obtain

$$\gamma \ll \omega_k \frac{N_c}{k_0^3}. \qquad (12)$$

From the comparison of conditions (9) and (12) one can see that they are equivalent to each other if $k_0^3 \geq N$, i.e., when the volume per one atom is larger than λ_0^3. According to (10), this means qualitatively that a single atom is fallen at many photons. Note, however, that condition (11) looses its meaning if the renormalization of the transition frequency is larger than the natural linewidth for a single atom, $|\varepsilon_b - \varepsilon_a - \omega_0| > \gamma$. In this case, it seems that γ should be regarded as a total natural linewidth with respect to the collisional broadening. The calculation of the corresponding contribution is made in [8].

In what follows, we assume that inequality (9) is fulfilled and $\omega \simeq \omega_k$. Denoting $\varepsilon = \varepsilon_a$ and re-denoting $\Omega_b \longrightarrow \Omega_b - \omega$, we have from the equations (7)

$$(\varepsilon - \Omega_a)(\varepsilon - \Omega_b) = |A|^2 \,.$$

Hence

$$\varepsilon = \frac{1}{2}\left[(\Omega_a + \Omega_b) + \eta\sqrt{\delta\omega^2 + 4|A|^2}\right], \eta = \pm 1. \qquad (13)$$

Here the quantity

$$\delta\omega = \Omega_a - \Omega_b = \omega - \omega_0 + \frac{\vec{k}^2}{2m}\left(\frac{N_b - N_a}{N_a + N_b}\right) + U_{aa}N_a - U_{bb}N_b \ + U_{ab}(N_b - N_a)$$

plays a role of the deviation from the full saturation. The relation between the amplitudes of the condensates is determined from (7)

$$b = \left[-\frac{\delta\omega}{2|A|} + \eta\sqrt{\left(\frac{\delta\omega}{2|A|}\right)^2 + 1}\right] a.$$

It follows from above that the system in the laser field is in one of the states with the fixed ratio of the condensate densities (at the given sign of product $\eta\delta\omega$)

$$N_b = \left[-\eta\frac{\delta\omega}{2|A|} + \sqrt{\left(\frac{\delta\omega}{2|A|}\right)^2 + 1}\right]^2 N_a. \qquad (14)$$

When $\delta\omega = 0$, the resonance field frequency equals $\omega = \omega_0 + \frac{1}{2}(U_{bb} - U_{aa})N$. Thus, if $U_{aa} \simeq U_{bb}$, it is natural to take difference $\omega - \omega_0$ as an external parameter characterizing the deviation from the saturation. Let us introduce the dimensionless variables

$$x = \frac{\omega - \omega_0}{2\,|A|},\ y = \frac{N_b - N_a}{N},\ z \simeq \frac{1}{2\,|A|}[\frac{k^2}{2m} + (U_{ab} - U_{aa})N] \tag{15}$$

and write equation (14) in the form

$$1 + y = \left(-\eta\,(x + zy) + \sqrt{(x + zy)^2 + 1}\right)^2 (1 - y). \tag{16}$$

Equation (16) is a quartic one with respect to $y\,(x)$. The solution relative to the inverse function $x\,(y)$ has a simple form

$$x = -(z + \frac{\eta}{\sqrt{1 - y^2}})y. \tag{17}$$

It follows from the above expression that there exists a symmetry with respect to the simultaneous substitution $x \longrightarrow -x$ and $y \longrightarrow -y$. This results from the symmetry of the Hamiltonian of the system with regard to substitution $\hat{a} \leftrightarrow \hat{b}$. In its turn, such symmetry between "up" and "down" is a result of the assumption about the smallness of γ. Note that there is no symmetry in the solutions with regard to the variation of the sign η. For the branch $\eta = +1$, quantity $y\,(x)$ is a one-to-one function of x and, for $\eta = -1$, function $y\,(x)$ at $z > 1$ within the interval

$$|x| < x_{\text{lim}} = \left(z^{2/3} - 1\right)^{3/2}$$

takes four values (see Fig.1). Accordingly, for the branch $\eta = -1$ at $z > 1$ there exists limiting values $y(\pm x_{\text{lim}}) = \mp\sqrt{1 - z^{-2/3}}$ with an anomalous S-shape of the line in comparison with the case $z < 1$.

For $x = 0$, both condensates have either the same density, $y = 0$, for any sign or

$$y = \pm\sqrt{1 - z^{-2}}, \tag{18}$$

at $\eta = -1$. Thus, the satellite states are possible if the coupling between atoms and field is sufficiently small compared with the recoil energy $\varepsilon_r = \frac{k^2}{2m}$ ($z > 1$, being $z \approx \frac{\varepsilon_r}{2|A|}$ for $U_{ab} \approx U_{aa}, U_{bb}$). If the coupling between atoms and field is strong, $z < 1$, there are solutions only in the vicinity of the centre ($y = -\frac{1}{z+\eta}x$) at $|x| \ll 1$ (i.e., in the case $|\varepsilon_b - \varepsilon_a - \omega_0| \leq \gamma \ll |A|$).

So, it is seen from the analysis of function $x\,(y)$ that (i) under the condition of saturation $|x| \ll 1$ there exist four stationary states, where in addition to the usual solutions $y \approx 0$ there are two satellite states in which populations of the ground and excited states are essentially different.

(ii) Beyond the region of saturation at $|x| > x_{\rm lim}$ there are two stationary states, and at $|x| < x_{\rm lim}$ there are four ones again. It is also interesting to trace the behavior $y(z)$ at $x = 0$. If $z > 1$, bifurcation, associated with a possibility of satellite states, appears in the plot $y(z)$ in addition to $y = 0$ (see Fig.2).

3. Combined oscillations of the condensates in the laser field

The stationary state obtained is unstable if the spectrum of elementary excitations of the system in this state has an imaginary part. To clarify a role of the interaction between atoms, we start from the case of an ideal gas. In addition, we will consider the question about the stability in the case of full saturation $\omega = \omega_0, N_a = N_b = N/2$ and in the case of a satellite state.

First, we find a set of equations determining spectrum of single-particle excitations of the condensates. Let us write equations (3) for the complex fields in the linear approximation in the amplitudes of noncondensate particles in the following representation (hereafter $\vec{q}$ is a quasiparticle momentum):

$$a_p = \bar{a}_p \exp\left(-i\varepsilon t\right), \quad \vec{p} = \vec{p_a} \pm \vec{q},$$

$$b_p = \bar{b}_p \exp\left(-i\varepsilon t - i\omega t\right), \quad \vec{p} = \vec{p_b} \pm \vec{q},$$

in which the system (3) takes a form of a closed system with constant coefficients (vector notations are omitted):

$$\begin{aligned} i\tfrac{\partial \bar{a}_{p_a+q}}{\partial t} &= (\tfrac{(p_a+q)^2}{2m} - \varepsilon + 2\mu_{aa} + U_{ab}N_b)\bar{a}_{p_a+q} + \mu_{aa}\bar{a}^{+}_{p_a-q} + (\mu_{ab}+A)\bar{b}_{p_b+q} + \mu_{ab}\bar{b}^{+}_{p_b-q}, \\ -i\tfrac{\partial \bar{a}^{+}_{p_a-q}}{\partial t} &= (\tfrac{(p_a-q)^2}{2m} - \varepsilon + 2\mu_{aa} + U_{ab}N_b)\bar{a}^{+}_{p_a-q} + \mu_{aa}\bar{a}_{p_a+q} + \mu_{ab}\bar{b}_{p_b+q} + (\mu_{ab}+A)\bar{b}^{+}_{p_b-q}, \\ i\tfrac{\partial \bar{b}_{p_b+q}}{\partial t} &= (\tfrac{(p_b+q)^2}{2m} - \varepsilon - (\omega-\omega_0) + 2\mu_{bb} + U_{ab}N_a)\bar{b}_{p_b+q} + \mu_{bb}\bar{b}^{+}_{p_b-q} + (\mu_{ab}+A)\bar{a}_{p_a+q} + \mu_{ab}\bar{a}^{+}_{p_a-q}, \\ -i\tfrac{\partial \bar{b}^{+}_{p_b-q}}{\partial t} &= (\tfrac{(p_b-q)^2}{2m} - \varepsilon - (\omega-\omega_0) + 2\mu_{bb} + U_{ab}N_a)\bar{b}^{+}_{p_b-q} + \mu_{bb}\bar{b}_{p_b+q} + \mu_{ab}\bar{a}_{p_a+q} + (\mu_{ab}+A)\bar{a}^{+}_{p_a-q}, \end{aligned} \tag{19}$$

where the notations $\mu_{ik} = U_{ik}\sqrt{N_i N_k}$ and $A = g^{*}_{k}c$ are used. The phase of laser field is chosen so that quantity A is real and positive.

In the case of an ideal gas, when all $U_{ik} = 0$, the equation for eigenvalues of the system (19), with involving (13) and (15), yields

$$E_{1,2} = -\frac{\vec{k}\vec{q}}{2m}y + \frac{q^2}{2m} - \frac{1}{2}\eta\sqrt{\delta\omega^2 + 4\left|A\right|^2} \pm \frac{1}{2}\sqrt{\left(\delta\omega + \frac{\vec{k}\vec{q}}{m}\right)^2 + 4\left|A\right|^2},$$

$$E_{3,4} = -\frac{\vec{k}\vec{q}}{2m}y - \frac{q^2}{2m} + \frac{1}{2}\eta\sqrt{\delta\omega^2 + 4\left|A\right|^2} \pm \frac{1}{2}\sqrt{\left(\delta\omega - \frac{\vec{k}\vec{q}}{m}\right)^2 + 4\left|A\right|^2}.$$

Here $\delta\omega = \omega - \omega_0 + \frac{k^2}{2m} y$. For the full saturation $(\delta\omega = 0, y = 0)$, we have

$$E_{1,2} = \left(\frac{q^2}{2m} - \eta\left|A\right|\right) \pm \sqrt{\left(\frac{\vec{k}\vec{q}}{2m}\right)^2 + \left|A\right|^2}, \quad E_{3,4} = -\frac{q^2}{2m} + \eta\left|A\right| \pm \sqrt{\left(\frac{\vec{k}\vec{q}}{2m}\right)^2 + \left|A\right|^2}.$$

Hence, for the given η there is a gap $2\left|A\right|$ for two modes in the excitation spectrum at $q = 0$. The gapless modes at $A = 0$ and $q \longrightarrow 0$ yield a sound dispersion law

$$E(q) \sim c_s q,$$

where $c_s = \frac{|\vec{k}\vec{q}|}{2mq}$ is the sound velocity. Far from the saturation $(|x| \gg 1, |y| \simeq 1)$ in (15) for the given η the two branches have a gap of about $|\omega - \omega_0|$ and the others are gapless as before.

Thus, in the approximation of an ideal gas the stationary condensate state is always stable in the sense mentioned above.

In the case of nonideal gas one can find an analytical expression for the spectrum of the system (19) under full saturation, assuming that $U_{aa} = U_{bb}$:

$$\begin{aligned} E_{\pm}^2 &= \left(P^2 - \mu^2\right) + A(A + 2\mu_{ab}) + Q^2 \pm 2\sqrt{(P^2 - \mu^2)Q^2 + (AP + \mu_{ab}(P - \mu))^2}, \\ P &= \frac{q^2}{2m} - \eta A + \mu, \quad Q = \frac{\vec{k}\vec{q}}{2m}, \quad \mu = \frac{1}{2} U_{aa} N, \quad \mu_{ab} = \frac{1}{2} U_{ab} N. \end{aligned} \tag{20}$$

First of all, we consider the long wave limit $q \longrightarrow 0$ and $E_{\pm}^2(q) \approx E_{\pm}^2(0) + \frac{q^2}{2m}\Delta E$ where

$$\begin{aligned} E_{\pm}^2(0) &= \xi^2 - \mu^2 + A\left(A + 2\mu_{ab}\right) \pm 2A\left|\xi - \eta\mu_{ab}\right|, \quad \xi = \mu - A\eta, \\ \Delta E &= 2\xi + \varepsilon_r \cos^2\theta \pm \left(\tfrac{(\xi^2 - \mu^2)}{A|\xi - \eta\mu_{ab}|}\varepsilon_r \cos^2\theta + 2A\left(A + \mu_{ab}\right) sign\left(\xi - \eta\mu_{ab}\right)\right). \end{aligned} \tag{21}$$

Here θ is the angle between vectors $\vec{k}$ and $\vec{q}$. The critical value, in the sense of stability, is E_{-}^2. In the case when the argument of modulus in (21) is positive $\mu > \eta\left(\mu_{ab} + A\right)$ we obtain

$$\begin{aligned} E_{-}^2(0) &= 2A(A - \mu + \mu_{ab})(1 + \eta), \\ \Delta E &= 2\left(\mu - \mu_{ab} - A\left(1 + \eta\right)\right) + \varepsilon_r \cos^2\theta\left[1 - \frac{(A - 2\eta\mu)}{\mu - \eta\left(\mu_{ab} + A\right)}\right]. \end{aligned}$$

It is seen that there is an instability in the state $\eta = 1$ at $\mu > \mu_{ab} + A$ and in the state $\eta = -1$

$$\Delta E = 2\left(\mu - \mu_{ab}\right)\left(1 - \frac{1}{2}\frac{\varepsilon_r \cos^2\theta}{\mu + \mu_{ab} + A}\right),$$

at $\mu < \mu_{ab}$. This coincides with the known condition [6] of instability of a binary Bose-gas mixture against spatial separation of its components. The instability at $\varepsilon_r \cos^2\theta > 2\left(\mu + \mu_{ab} + A\right)$ means that for the strong recoil the instability develops at first in the direction of the recoil momentum.

The corresponding plot of the spectrum (20) at $\cos^2\theta = 1$ is given in Fig.3. The opposite case $\mu < \eta\left(\mu_{ab} + A\right)$ has a sense only for $\eta = 1$ and does not result in an instability

$$E_-^2(0) = 0, \quad \Delta E = 2\left(\mu + \mu_{ab}\right)\left(1 + \frac{1}{2}\frac{\varepsilon_r\cos^2\theta}{A + \mu_{ab} - \mu}\right) > 0.$$

Let us now consider the case of transverse excitations with respect to vector $\vec{k}$ when $Q = 0$

$$E_\pm^2 = \delta^2 + 2\mu\delta + A(A + 2\mu_{ab}) \pm 2\left|A\left(\delta + \mu\right) + \mu_{ab}\delta\right|, \quad \delta = \frac{q^2}{2m} - \eta A. \tag{22}$$

For E_-^2 at $\left(A + \mu_{ab}\right)\left(\frac{q^2}{2m} - \eta A\right) + A\mu > 0$, we get

$$E_-^2 = \left(\frac{q^2}{2m} - A\left(1 + \eta\right)\right)\left(\frac{q^2}{2m} - A\left(1 + \eta\right) + 2\left(\mu - \mu_{ab}\right)\right).$$

It is seen that in the state $\eta = -1$ an instability appears for the condition obtained above $\mu < \mu_{ab}$. In the state $\eta = 1$ there is a region of instability $2A - 2\left(\mu - \mu_{ab}\right) < \frac{q^2}{2m} < 2A$ with the centre at the point $\frac{q_{\min}^2}{2m} = 2A - \left(\mu - \mu_{ab}\right)$ where $E_-^2(q_{\min}) = -\left(\mu - \mu_{ab}\right)^2$. In addition, for $q = 0$ in this state there appears an instability if $\mu > A + \mu_{ab}$ (however, it is always $E_-^2(q_{\min}) < E_-^2(0)$). The typical plot of the spectrum for this case (20) is given in Fig.4.

In the case $\left(A + \mu_{ab}\right)\left(\frac{q^2}{2m} - \eta A\right) + A\mu < 0$ we have

$$E_-^2 = \left(\frac{q^2}{2m} + A\left(1 - \eta\right)\right)\left(\frac{q^2}{2m} + A\left(1 - \eta\right) + 2\left(\mu + \mu_{ab}\right)\right),$$

whence it follows that the both states $\eta = \pm 1$ are stable.

Finally, let us consider the stability of satellites at $U_{aa} = U_{bb}$. Far from the saturation ($|\omega - \omega_0| \gg A,\ \mu,\ \mu_{ab}$) one can neglect the nonideality of the gas and, as we convinced above, the condensates are stable. In the region of saturation $|x| \ll 1$, the satellites exist only in the state with $\eta = -1$. For simplicity, we restrict ourselves with the case when the density of one of condensates is much less than that of the other. For definiteness, $N_b \ll N_a$ ($y \simeq -1$). It turns out that one can derive an analytical expression for the spectrum of elementary excitations in the approximation of heavy atoms and strong atom-atom coupling compared with the recoil, putting formally $k \ll q$. Note that in these conditions it is not allowed to employ the long wave limit $q \longrightarrow 0$.

$$E_\pm^2 = \frac{1}{2}\left(W^2 + V^2 + 2A^2 - \mu^2 \pm \sqrt{\left(W^2 - V^2 - \mu^2\right)^2 + 4A^2\left(\left(W + V\right)^2 - \mu^2\right)}\right), \tag{23}$$

$$W = \frac{q^2}{2m} + \frac{1}{2}\left(3\mu - \mu_{ab}\right) + \sqrt{\frac{1}{4}\left(\mu - \mu_{ab}\right)^2 + A^2},$$

$$V = \frac{q^2}{2m} - \frac{1}{2}\left(\mu - \mu_{ab}\right) + \sqrt{\frac{1}{4}\left(\mu - \mu_{ab}\right)^2 + A^2},$$

where $\mu = U_{aa}N$, $\mu_{ab} = U_{ab}N$. Rewriting expression (23) in the form

$$E_{\pm}^{2} = \frac{1}{2}\left(W^{2}+V^{2}+2A^{2}-\mu^{2} \pm \sqrt{\left(W^{2}+V^{2}+2A^{2}-\mu^{2}\right)^{2}-4\left(A^{2}-WV\right)^{2}+4\mu^{2}V^{2}}\right),$$

one can straightforwardly indicate the region of possible negative values E_{-}^{2},

$$-4\left(A^{2}-WV\right)^{2}+4\mu^{2}V^{2}>0.$$

Hence

$$\mu V > \left(WV - A^{2}\right). \tag{24}$$

Note that the condition (24) never holds true at $\mu \simeq \mu_{ab}$ and the satellite state is stable.

4. Conclusion

The main result of the work is the determination of a stationary solution of the equations of motion for the united coherent system of two-level atom Bose condensates and laser field of high intensity. In addition, the dependence of the condensate densities on the Rabi frequency and deviation from the full saturation are found. The analysis is performed for the stability of the system against appearance of the imaginary part in the spectrum of elementary excitations. This results in an exponential increase of oscillations, in the heating of the system and decay of the condensates, or in the transition of the system into one of the stable states. The question on evolution of the system after the decay of an unstable stationary state is a subject of separate study.

The condition of applicability of the theory in temperature is given by

$$T \ll T_c \sim \frac{\hbar^{2}}{m} n^{2/3}, \quad n = \min\left(N_a, N_b\right).$$

Note that in the satellite state, when the condensate densities differ significantly, the condensate of the lower density at $T > T_c$ converts into noncondensate particles which are additional with respect to the condensate of higher density. The question whether this condensate will be destroyed with adding extrinsic noncondensate particles represents an independent interest.

The authors are grateful to Yu. Kagan and S. N. Burmistrov for valuable comments. The work is supported by the grants of Russian Foundation of Basic Research and INTAS.

---

[1] M. H. Anderson, J. R. Ensher, M. R. Matthews *et al.*, Science **269**, 198 (1995); K. B. Davis, M. -O. Mewes, M. R. Andrews *et al.*, Phys. Rev. Lett. **75**, 3969 (1995); C. C. Bradley, C. A. Sackett, and R.

G. Hulet, Phys. Rev. Lett. **78**, 985 (1997); D. G. Fried, Th. C. Killian, L. Willmann *et al.*, Phys. Rev. Lett. **81**, 3811 (1998).

[2] D. S. Hall, M. R. Matthews, J. R. Ensher *et al.*, Phys. Rev. Lett. **81**, 1539 (1998).

[3] Tin-Lin Ho and V. B. Shenoy, Phys. Rev. Lett. **77**, 3276 (1996); M. Trippenbach, K. Goral, K. Rzazewski *et al.*, J. Phys. B **33**, 4017 (2000).

[4] Th. Busch, J. I. Cirac, V. M. Perez-Garcia and P. Zoller, Phys. Rev. A **56**, 2978 (1997); R. Graham and D. Walls, Phys. Rev. A **57**, 484 (1998).

[5] H. Pu and N. P. Bigelow, Phys. Rev. Lett. **80**, 1134 (1998).

[6] P. Ao and S. T. Chui, Phys. Rev. A **58**, 4836 (1998); E. Timmermans, Phys. Rev. Lett. **81**, 5718 (1998); H. Shi, Wei-Mou Zheng and Siu-Tat Chui, Phys. Rev. A **61**, 063613 (1998); R. A. Barankov, Phys. Rev. A **66**, 013612 (2002).

[7] J. Ruostekoski, M. J. Collett, R. Graham and D. Walls, Phys. Rev. A **57**, 511 (1998).

[8] M. O. Oktel, L. S. Levitov, Phys. Rev. Lett. **83**, 6 (1999).

Figure Captions

Fig.1. The dependence of the relative difference in populations, y, versus deviation x from the full saturation.

Fig.2. The dependence of the relative difference in populations, y, versus parameter z (see (15)).

Fig.3. The example of unstable spectrum in the long wave limit for the state $\eta = -1$ under full saturation at $\varepsilon_r > 2\,(\mu + \mu_{ab} + A) = 0.43\varepsilon_r$, and $\mu : \mu_{ab} : A = 1 : 0.5 : 20$. The scale on the vertical axis for the curve $\eta = 1$ is reduced by a factor of 40.

Fig.4. An example of the unstable spectrum for the transverse excitations in the state $\eta = 1$ under full saturaion for $Q = 0$, $A = 2.5\varepsilon_r$, $\mu : \mu_{ab} : A = 3 : 1 : 5$. The scale of the vertical axis of curve $\eta = -1$ is reduced by a factor of 4. The cusp in the curve E_{-}^{2}, $\eta = 1$, corresponds to zero modulus in (22).



Figure 1:



Figure 2:


x 10^{−5}
6
5
4
3
2
1
0
−1
−2
−3
E_−^2/ε_r^2
η = 1
η = −1
0
0.01
0.02
0.03
0.04
0.05
0.06
0.07
0.08
0.09
0.1
q/k

Figure 3:



Figure 4: